

English SIG Code Guide

The Patient Takeaway that you provide as a follow-up to a Comprehensive Medication Review (CMR) is an important resource for your patients. To ensure a patient can easily understand the information, please use patient-friendly language by avoiding abbreviations and unfamiliar vocabulary.

To save you time, the Connect platform translates SIG codes into patient-friendly language. For example, “BID AC” (note the space between “BID” and “AC”) translates to “TWICE DAILY BEFORE A MEAL” and “1TBIDAC” translates to “TAKE 1 TABLET DAILY BEFORE MEALS”.

NOTE: You should always review the directions to ensure everything has translated correctly before printing a takeaway for the patient.

Where can I use SIG codes in Connect?

SIG codes work in the Medication List on the Patient Profile and within CMR documentation.

How do I use SIG codes?

1. Enter the appropriate SIG code(s) for each medication in the Directions field.
2. If you move the cursor outside the text field or press the TAB button to move to the next field, the SIG code translation will appear.

Best practices for using SIGs

- Adding a space between individual SIG codes lets you create any combination of codes.
- Some combination SIG codes do not require a space between them.
- Free text may be incorporated with SIG codes as long as there is a space between the text and SIG code. For example, “APPLY TO THE AFFECTED AREA” can be entered as “APPLY AA” (note the space between “APPLY” and “AA”).
- SIG codes are not case sensitive.
- Some measurements, such as MEQ, MG, ML and MLS, remain in shortened form.

English SIG Code Translations

Common SIG codes

Oral - Capsules

SIG code	Description
1CBID	TAKE 1 CAPSULE BY MOUTH TWICE DAILY
1CQAM	TAKE 1 CAPSULE BY MOUTH EVERY MORNING
1CQD	TAKE 1 CAPSULE BY MOUTH ONCE DAILY

Oral - Tablets

SIG code	Description
1TBID	TAKE 1 TABLET BY MOUTH TWICE DAILY
1TQAM	TAKE 1 TABLET BY MOUTH EVERY MORNING
1TQD	TAKE 1 TABLET BY MOUTH ONCE DAILY
1TQPM	TAKE 1 TABLET BY MOUTH EVERY EVENING

Oral - Inhalation

SIG code	Description
1INHQD	USE 1 INHALATION BY MOUTH ONCE DAILY
1PQ4-6PRN	INHALE 1 PUFF BY MOUTH EVERY 4 TO 6 HOURS AS NEEDED
2PBID	INHALE 2 PUFFS BY MOUTH TWICE

Actions

SIG code	Description
CRU	CRUSH
CSW	CHEW AND SWALLOW
CW	CHEW
DK	DRINK
INH	INHALE
INJ	INJECT
INS	INSTILL

English SIG Code Translations

Actions continued

SIG code	Description
SAS	SWISH AND SWALLOW
SHW	SHAKE WELL
SP	SPRAY
SSP	SWISH AND SPIT
TK	TAKE
UNW	UNWRAP

Other

SIG code	Description
COAG	AS DIRECTED BY ANTICOAGULATION CLINIC
DAPP	DENTAL APPOINTMENT
DTI	DOUBLE OR TRIPLE FOR ILLNESS
FPD	FOLLOW PACKAGE DIRECTIONS
HA	HEADACHE
MCD	MAY CAUSE DROWSINESS
NA	NAUSEA
NER	FOR NERVES
NM3	NO MORE THAN 3,000 MG ACETAMINOPHEN DAILY
NM4	NO MORE THAN 4,000 MG ACETAMINOPHEN DAILY
NTE	NOT TO EXCEED
NTEAPAP	NOT TO EXCEED 3000 MG IN 24 HOURS
NV	FOR NAUSEA AND VOMITING
P	PAIN
UAD	USE AS DIRECTED
UD	AS DIRECTED
UDMD	AS DIRECTED BY YOUR DOCTOR
VOM	VOMITING

English SIG Code Translations

Dosage form

SIG code	Description
1C	1 CAPSULE
1GU	1 GUMMY
1T	1 TABLET
1V	1 VIAL
2C	2 CAPSULES
2GU	2 GUMMIES
2T	2 TABLETS
2V	2 VIALS
3C	3 CAPSULES
3T	3 TABLETS
4C	4 CAPSULES
4T	4 TABLETS
C	CAPSULE
CS	CAPSULES
CW	CHEW
CWS	CHEWS
GTT	DROP
GTTS	DROPS
GU	GUMMY
LQ	LIQUID
NEB	NEBULIZER
OZ	OUNCE
OZS	OUNCES
PF	PUFF
PFS	PUFFS
SP	SPRAY
SPS	SUPPOSITORIES
SPY	SPRAYS
SSI	SLIDING SCALE INSULIN

English SIG Code Translations

Dosage form continued

SIG code	Description
SUP	SUPPOSITORY
SYR	SYRINGE
T	TABLET
TS	TABLETS

Frequency

SIG code	Description
30MACAM	30 MINUTES BEFORE MORNING MEAL
60MACAM	60 MINUTES BEFORE MORNING MEAL
AC	BEFORE A MEAL
AF	AFTER
AFT	IN THE AFTERNOON
AM	MORNING
B	BEFORE
BID	TWICE A DAY
BIW	TWICE A WEEK
D	DAILY
FEV	AS NEEDED FOR FEVER
FEVP	AS NEEDED FOR FEVER AND PAIN
FID	5 TIMES DAILY
H	HOURS
HS	AT BEDTIME
IMM	IMMEDIATELY
MWF	ON MONDAYS, WEDNESDAYS, AND FRIDAYS
OES	ON EMPTY STOMACH
PC	AFTER A MEAL
PM	EVENING
PP	AS NEEDED FOR PAIN

English SIG Code Translations

Frequency continued

SIG code	Description
PRA	AS NEEDED FOR ANXIETY
PRF	AS NEEDED FOR
PRN	AS NEEDED
PTB	PER TUBE
Q	EVERY
Q1M	EVERY MONTH
Q1W	EVERY WEEK
Q2W	EVERY 2 WEEKS
Q3D	EVERY 3 DAYS
Q3M	EVERY 3 MONTHS
Q4-6	EVERY 4-6 HOURS
Q4-6H	EVERY 4-6 HOURS
Q4-6HPRN	EVERY 4-6 HOURS AS NEEDED
Q4-6PRN	EVERY 4-6 HOURS AS NEEDED
Q4H	EVERY 4 HOURS
Q4W	EVERY 4 WEEKS
Q6H	EVERY 6 HOURS
Q6M	EVERY 6 MONTHS
Q7D	EVERY 7 DAYS
Q8H	EVERY 8 HOURS
Q12H	EVERY 12 HOURS
QAM	EVERY MORNING
QD	ONE TIME A DAY
QHS	EVERY NIGHT AT BEDTIME
QID	4 TIMES DAILY
QN	EVERY NIGHT
QOD	EVERY OTHER DAY
QOW	EVERY OTHER WEEK
QPM	EVERY EVENING

English SIG Code Translations

Frequency continued

SIG code	Description
TAT	UNTIL ALL TAKEN
TID	THREE TIMES DAILY
TIW	THREE TIMES A WEEK
TTSS	ON TUESDAY, THURSDAY, SATURDAY, AND SUNDAY
WA	WHILE AWAKE
WC	WITH MEALS
WF	WITH FOOD
WK	WEEK
XD	TIMES DAILY
XWK	TIMES WEEKLY

Measurements

SIG code	Description
CC	CUBIC CENTIMETER
CFL	CAPFUL
GM	GM
L	L
MCG	MCG
MEQ	MEQ
MG	MG
ML	ML
MLS	MLS
OZ	OUNCE
OZS	OUNCE
SS	ONE-HALF
TES	TEASPOONFUL (5 ML)

English SIG Code Translations

Measurements continued

SIG code	Description
TEA	TEASPOON
TBL	TABLESPOON
TBLS	TABLESPOONFUL (15 ML)

Route of administration

SIG code	Description
AA	TO THE AFFECTED AREA
AD	RIGHT EAR
AU	BOTH EARS
EXT	EXTERNALLY
IAD	IN THE RIGHT EAR
IAS	IN THE LEFT EAR
IAU	IN BOTH EARS
IEN	IN EACH NOSTRIL
IOD	IN THE RIGHT EYE
IOS	IN THE LEFT EYE
IOU	IN EACH EYE
IM	IN THE MUSCLE
IV	INTO THE VEIN
NOS	NOSTRIL
OD	RIGHT EYE
OS	LEFT EYE
OU	EACH EYE
PO	BY MOUTH
REC	RECTALLY
SC	SUBCUTANEOUS OR UNDER THE SKIN
SL	SUBLINGUAL

English SIG Code Translations

Route of administration continued

SIG code	Description
SQ	UNDER THE SKIN
UNT	UNDER THE TONGUE
VAG	IN THE VAGINA

Combination SIG codes - Injectable

SIG code	Description
BASINS	INJECT UNITS OF INSULIN UNDER THE SKIN ONCE DAILY AS DIRECTED BY YOUR PRESCRIBER
BOLINS	INJECT UNITS OF INSULIN UNDER THE SKIN BEFORE MEALS AS DIRECTED BY YOUR PRESCRIBER
EPIPENSIG	INJECT THE CONTENTS OF 1 PEN INTO THE OUTER THIGH IF NEEDED FOR A SEVERE ALLERGIC REACTION. MAY REPEAT DOSE EVERY 5 TO 15 MINUTES AS NEEDED. CALL 911 IF NO IMPROVEMENT.
GLUCAGONSIG	MIX CONTENTS OF SYRINGE AND VIAL, THEN DRAW UP SOLUTION INTO THE SYRINGE. INJECT SOLUTION INTO THE SKIN AS NEEDED FOR LOW BLOOD SUGAR.
ISSI	INJECT INSULIN UNDER THE SKIN BEFORE MEALS ACCORDING TO SLIDING SCALE PROVIDED BY YOUR PRESCRIBER
NALOXINJ	PRESS THE AUTOINJECTOR (2 MG) INTO OUTER PORTION OF THIGH AND HOLD IN PLACE FOR 5 SECONDS. SEEK EMERGENCY MEDICAL ASSISTANCE AFTER USE. MAY GIVE ANOTHER INJECTION IN 2 TO 3 MINUTES IF NO RESPONSE.
PROLIASIG	INJECT 60 MG UNDER THE SKIN EVERY 6 MONTHS
SSIUMD	ACCORDING TO SLIDING SCALE PROVIDED BY YOUR PRESCRIBER

Combination SIG codes - Nasal

SIG code	Description
1INHENBID	USE 1 INHALATION IN EACH NOSTRIL TWICE DAILY
1INHENQD	USE 1 INHALATION IN EACH NOSTRIL ONCE DAILY

English SIG Code Translations

Combination SIG codes - Nasal continued

SIG code	Description
1SPENBID	USE 1 SPRAY IN EACH NOSTRIL TWICE DAILY
1SPENHS	USE 1 SPRAY IN EACH NOSTRIL AT BEDTIME
1SPENQAM	USE 1 SPRAY IN EACH NOSTRIL EVERY MORNING
1SPENQD	USE 1 SPRAY IN EACH NOSTRIL ONCE DAILY
1-2SPENBID	USE 1 OR 2 SPRAYS IN EACH NOSTRIL TWICE DAILY
1-2SPENQD	USE 1 OR 2 SPRAYS IN EACH NOSTRIL ONCE DAILY
2INHENBID	USE 2 INHALATIONS IN EACH NOSTRIL TWICE DAILY
2INHENQD	USE 2 INHALATIONS IN EACH NOSTRIL ONCE DAILY
2SPENBID	USE 2 SPRAYS IN EACH NOSTRIL TWICE DAILY
2SPENHS	USE 2 SPRAYS IN EACH NOSTRIL AT BEDTIME
2SPENQAM	USE 2 SPRAYS IN EACH NOSTRIL EVERY MORNING
2SPENQD	USE 2 SPRAYS IN EACH NOSTRIL ONCE DAILY
BAQSIMISIG	USE 1 SPRAY (3 MG) IN ONE NOSTRIL AS NEEDED FOR SYMPTOMS OF LOW BLOOD SUGAR. IF NO RESPONSE, MAY REPEAT DOSE IN ALTERNATE NOSTRIL IN 15 MINUTES USING A NEW DEVICE.
NALOXSP	USE 1 SPRAY (4 MG) IN ONE NOSTRIL FOR SYMPTOMS OF OPIOID OVERDOSE AND SEEK EMERGENCY MEDICAL ASSISTANCE. REPEAT DOSE IN ALTERNATE NOSTRIL IN 2 TO 3 MINUTES IF NO RESPONSE.

Combination SIG codes - Ophthalmic

SIG code	Description
1GODBID	USE 1 DROP IN RIGHT EYE TWICE A DAY
1GODHS	USE 1 DROP IN RIGHT EYE AT BEDTIME
1GODQD	USE 1 DROP IN RIGHT EYE ONCE DAILY
1GOSBID	USE 1 DROP IN LEFT EYE TWICE A DAY
1GOSHS	USE 1 DROP IN LEFT EYE AT BEDTIME
1GOSQD	USE 1 DROP IN LEFT EYE ONCE DAILY
1GOUAM	USE 1 DROP IN EACH EYE IN THE MORNING

English SIG Code Translations

Combination SIG codes - Ophthalmic continued

SIG code	Description
1GOUBID	USE 1 DROP IN EACH EYE TWICE DAILY
1GOUHS	USE 1 DROP IN EACH EYE AT BEDTIME
1GOUQD	USE 1 DROP IN EACH EYE ONCE DAILY
1GOUTID	USE ONE DROP IN EACH EYE THREE TIMES DAILY

Combination SIG codes - Oral Inhalation

SIG code	Description
1INHBID	USE 1 INHALATION BY MOUTH TWICE DAILY
1INHQAM	USE 1 INHALATION BY MOUTH EVERY MORNING
1INHQD	USE 1 INHALATION BY MOUTH ONCE DAILY
1PBID	INHALE 1 PUFF BY MOUTH TWICE DAILY
1PQ4-6PRN	INHALE 1 PUFF BY MOUTH EVERY 4 TO 6 HOURS AS NEEDED
1PQD	INHALE 1 PUFF BY MOUTH ONCE DAILY
1-2PBID	INHALE 1 TO 2 PUFFS BY MOUTH TWICE DAILY
1-2PQ4	INHALE 1 TO 2 PUFFS BY MOUTH EVERY 4 HOURS
1-2PQ4-6PRN	INHALE 1 TO 2 PUFFS BY MOUTH EVERY 4 TO 6 HOURS AS NEEDED
1-2PQ4PRN	INHALE 1 TO 2 PUFFS BY MOUTH EVERY 4 HOURS AS NEEDED
1-2PQ6	INHALE 1 TO 2 PUFFS BY MOUTH EVERY 6 HOURS
1-2PQ6PRN	INHALE 1 TO 2 PUFFS BY MOUTH EVERY 6 HOURS AS NEEDED
2INHBID	USE 2 INHALATIONS BY MOUTH TWICE DAILY
2INHQD	USE 2 INHALATIONS BY MOUTH ONCE DAILY
2INHQAM	USE 2 INHALATIONS BY MOUTH EVERY MORNING
2PBID	INHALE 2 PUFFS BY MOUTH TWICE DAILY
2PQ4-6PRN	INHALE 2 PUFFS BY MOUTH EVERY 4-6 HOURS AS NEEDED
2PQ46H	INHALE 2 PUFFS BY MOUTH INTO THE LUNGS EVERY 4 TO 6 HOURS
2PQD	INHALE TWO PUFFS BY MOUTH ONCE DAILY
COMBIRESP	USE 1 INHALATION BY MOUTH 4 TIMES DAILY. TWIST DEVICE UNTIL CLICK, REMOVE CAP, AND PLACE MOUTHPIECE IN MOUTH. PUSH BUTTON, TAKE A DEEP BREATH, AND HOLD FOR 5 TO 10 SECONDS.

English SIG Code Translations

Combination SIG codes - Oral Inhalation continued

SIG code	Description
FLEXHALERSIG	PLACE THE MOUTHPIECE IN YOUR MOUTH, CLOSE YOUR LIPS AROUND IT, AND INHALE DEEPLY. USE 2 INHALATIONS BY MOUTH TWICE DAILY
HANDIHALERSIG	INHALE THE CONTENTS OF 1 CAPSULE VIA HANDIHALER BY MOUTH ONCE DAILY
RESPIMATSIG	USE 2 INHALATIONS BY MOUTH ONCE DAILY. TWIST DEVICE UNTIL CLICK, REMOVE CAP, PLACE MOUTHPIECE IN YOUR MOUTH. PUSH BUTTON, TAKE A SLOW DEEP BREATH. HOLD FOR 5 TO 10 SECONDS.
VVN	USE NEBULIZER TO INHALE THE CONTENTS OF 1 VIAL BY MOUTH
VVNBID	USE NEBULIZER TO INHALE THE CONTENTS OF 1 VIAL BY MOUTH TWICE A DAY
VVNQ4H	USE NEBULIZER TO INHALE THE CONTENTS OF 1 VIAL BY MOUTH EVERY 4 HOURS
VVNQ4-6HPRN	USE NEBULIZER TO INHALE THE CONTENTS OF 1 VIAL BY MOUTH EVERY 4 TO 6 HOURS AS NEEDED
VVNQD	USE NEBULIZER TO INHALE THE CONTENTS OF 1 VIAL BY MOUTH ONCE DAILY
VVNQID	USE NEBULIZER TO INHALE THE CONTENTS OF 1 VIAL BY MOUTH FOUR TIMES A DAY
VVNTID	USE NEBULIZER TO INHALE THE CONTENTS OF 1 VIAL BY MOUTH THREE TIMES A DAY

English SIG Code Translations

Combination SIG codes - Oral

SIG code	Description
.5TAC	TAKE 1/2 TABLET BY MOUTH BEFORE MEALS
.5TACHS	TAKE 1/2 TABLET BY MOUTH BEFORE MEALS AND AT BEDTIME
.5TAM	TAKE 1/2 TABLET BY MOUTH DAILY IN THE MORNING
.5TBID	TAKE 1/2 TABLET BY MOUTH TWICE DAILY
.5TBIDPRN	TAKE 1/2 TABLET BY MOUTH TWICE DAILY AS NEEDED
.5THS	TAKE 1/2 TABLET BY MOUTH AT BEDTIME
.5THSPRN	TAKE 1/2 TABLET BY MOUTH AT BEDTIME AS NEEDED
.5TPM	TAKE 1/2 TABLET BY MOUTH IN THE EVENING
.5TPOBID	TAKE 1/2 TABLET BY MOUTH TWICE DAILY
.5TPOHS	TAKE 1/2 TABLET BY MOUTH AT BEDTIME
.5TPOQD	TAKE 1/2 TABLET BY MOUTH ONCE DAILY
.5TPRN	TAKE 1/2 TABLET BY MOUTH AS NEEDED
.5TQ4-6PRN	TAKE 1/2 TABLET BY MOUTH EVERY 4 TO 6 HOURS AS NEEDED
.5TQ4PRN	TAKE 1/2 TABLET BY MOUTH EVERY 4 HOURS AS NEEDED
.5TQ6-8PRN	TAKE 1/2 TABLET BY MOUTH EVERY 6 TO 8 HOURS AS NEEDED
.5TQ6PRN	TAKE 1/2 TABLET BY MOUTH EVERY 6 HOURS AS NEEDED
.5TQ8	TAKE 1/2 TABLET BY MOUTH EVERY 8 HOURS
.5TQAM	TAKE 1/2 TABLET BY MOUTH EVERY MORNING
.5TQD	TAKE 1/2 TABLET BY MOUTH ONCE DAILY
.5TQDAM	TAKE 1/2 TABLET BY MOUTH ONCE DAILY IN THE MORNING
.5TQDHS	TAKE 1/2 TABLET BY MOUTH DAILY AT BEDTIME
.5TQDPM	TAKE 1/2 TABLET BY MOUTH ONCE DAILY IN THE EVENING
.5TQDPRN	TAKE 1/2 TABLET BY MOUTH DAILY AS NEEDED
.5TQHS	TAKE 1/2 TABLET BY MOUTH EVERY NIGHT AT BEDTIME
.5TQID	TAKE 1/2 TABLET BY MOUTH 4 TIMES A DAY
.5TQIDPRN	TAKE 1/2 TABLET BY MOUTH 4 TIMES A DAY AS NEEDED
.5TQOD	TAKE 1/2 TABLET BY MOUTH EVERY OTHER DAY
.5TTID	TAKE 1/2 TABLET BY MOUTH 3 TIMES A DAY
.5TTIDAC	TAKE 1/2 TABLET BY MOUTH 3 TIMES A DAY BEFORE MEALS

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
.5TTIDPRN	TAKE 1/2 TABLET BY MOUTH 3 TIMES A DAY AS NEEDED
.5-1TPOQD	TAKE 1/2 TO 1 TABLET BY MOUTH ONCE DAILY
.5-1TQD	TAKE 1/2 TO 1 TABLET BY MOUTH ONCE DAILY
.5-1TQDPRN	TAKE 1/2 TO 1 TABLET BY MOUTH ONCE DAILY AS NEEDED
.5-1TQID	TAKE 1/2 TO 1 TABLET BY MOUTH 4 TIMES A DAY
.5-1TTID	TAKE 1/2 TO 1 TABLET BY MOUTH 3 TIMES A DAY
.5-1TTIDPRN	TAKE 1/2 TO 1 TABLET BY MOUTH 3 TIMES A DAY AS NEEDED
1CAC	TAKE 1 CAPSULE BY MOUTH BEFORE MEALS
1CACHS	TAKE 1 CAPSULE BY MOUTH BEFORE MEALS AND AT BEDTIME
1CAM	TAKE 1 CAPSULE BY MOUTH IN THE MORNING
1CAM1CPM	TAKE 1 CAPSULE BY MOUTH IN THE MORNING AND TAKE 1 CAPSULE BY MOUTH IN THE EVENING
1CAMHS	TAKE 1 CAPSULE BY MOUTH IN THE MORNING AND AT BEDTIME
1CAMP	TAKE 1 CAPSULE BY MOUTH IN THE MORNING AND EVENING
1CBID	TAKE 1 CAPSULE BY MOUTH TWICE DAILY
1CBIDAC	TAKE 1 CAPSULE BY MOUTH TWICE DAILY BEFORE MEALS
1CBIDCF	TAKE 1 CAPSULE BY MOUTH TWICE DAILY WITH FOOD
1CBIDPRN	TAKE 1 CAPSULE BY MOUTH TWICE DAILY AS NEEDED
1CHS	TAKE 1 CAPSULE BY MOUTH AT BEDTIME
1CHSPRN	TAKE 1 CAPSULE BY MOUTH AT BEDTIME AS NEEDED
1CPM	TAKE 1 CAPSULE BY MOUTH IN THE EVENING
1CPMPRN	TAKE 1 CAPSULE BY MOUTH IN THE EVENING AS NEEDED
1CPOBID	TAKE 1 CAPSULE BY MOUTH TWICE DAILY
1CPOBIDPRN	TAKE 1 CAPSULE BY MOUTH TWICE DAILY AS NEEDED
1CPOHS	TAKE 1 CAPSULE BY MOUTH AT BEDTIME
1CPOQAM	TAKE 1 CAPSULE BY MOUTH EVERY MORNING
1CPOQAMAC	TAKE 1 CAPSULE BY MOUTH EVERY MORNING BEFORE MEALS
1CPOQD	TAKE 1 CAPSULE BY MOUTH ONCE DAILY
1CPOQDPRN	TAKE 1 CAPSULE BY MOUTH ONCE DAILY AS NEEDED

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
1CPOQHS	TAKE 1 CAPSULE BY MOUTH EVERY DAY AT BEDTIME
1CPOQID	TAKE 1 CAPSULE BY MOUTH 4 TIMES A DAY
1CPOQIDPRN	TAKE 1 CAPSULE BY MOUTH 4 TIMES A DAY AS NEEDED
1CPOQM	TAKE 1 CAPSULE BY MOUTH ONCE MONTHLY
1CPOQOD	TAKE 1 CAPSULE BY MOUTH EVERY OTHER DAY
1CPOQPM	TAKE 1 CAPSULE BY MOUTH EVERY EVENING
1CPOQWK	TAKE 1 CAPSULE BY MOUTH ONCE WEEKLY
1CPOTID	TAKE 1 CAPSULE BY MOUTH 3 TIMES A DAY
1CPOTIDPRN	TAKE 1 CAPSULE BY MOUTH 3 TIMES A DAY AS NEEDED
1CQ12H	TAKE 1 CAPSULE BY MOUTH EVERY 12 HOURS
1CQ3-4PRN	TAKE 1 CAPSULE BY MOUTH EVERY 3 TO 4 HOURS AS NEEDED
1CQ3-4QDPRN	TAKE 1 CAPSULE BY MOUTH EVERY 3 TO 4 HOURS DAILY AS NEEDED
1CQ3PRN	TAKE 1 CAPSULE BY MOUTH EVERY 3 HOURS AS NEEDED
1CQ4	TAKE 1 CAPSULE BY MOUTH EVERY 4 HOURS
1CQ4-6	TAKE 1 CAPSULE BY MOUTH EVERY 4 TO 6 HOURS
1CQ4-6PRN	TAKE 1 CAPSULE BY MOUTH EVERY 4 TO 6 HOURS AS NEEDED
1CQ4PRN	TAKE 1 CAPSULE BY MOUTH EVERY 4 HOURS AS NEEDED
1CQ6	TAKE 1 CAPSULE BY MOUTH EVERY 6 HOURS
1CQ6-8	TAKE 1 CAPSULE BY MOUTH EVERY 6 TO 8 HOURS
1CQ6-8PRN	TAKE 1 CAPSULE BY MOUTH EVERY 6 TO 8 HOURS AS NEEDED
1CQ6CF	TAKE 1 CAPSULE BY MOUTH EVERY 6 HOURS WITH FOOD
1CQ6PRN	TAKE 1 CAPSULE BY MOUTH EVERY 6 HOURS AS NEEDED
1CQ8	TAKE 1 CAPSULE BY MOUTH EVERY 8 HOURS
1CQ8CF	TAKE 1 CAPSULE BY MOUTH EVERY 8 HOURS WITH FOOD
1CQ8PRN	TAKE 1 CAPSULE BY MOUTH EVERY 8 HOURS AS NEEDED
1CQAM	TAKE 1 CAPSULE BY MOUTH EVERY MORNING
1CQAMAC	TAKE 1 CAPSULE BY MOUTH EVERY MORNING BEFORE MEALS
1CQAMPRN	TAKE 1 CAPSULE BY MOUTH EVERY MORNING AS NEEDED

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
1CQD	TAKE 1 CAPSULE BY MOUTH ONCE DAILY
1CQDAM	TAKE 1 CAPSULE BY MOUTH ONCE DAILY IN THE MORNING
1CQDAMCF	TAKE 1 CAPSULE BY MOUTH ONCE DAILY IN THE MORNING WITH FOOD
1CQDCF	TAKE 1 CAPSULE BY MOUTH ONCE DAILY WITH FOOD
1CQDHS	TAKE 1 CAPSULE BY MOUTH ONCE DAILY AT BEDTIME
1CQDHSPRN	TAKE 1 CAPSULE BY MOUTH ONCE DAILY AT BEDTIME AS NEEDED
1CQDP	TAKE 1 CAPSULE BY MOUTH ONCE DAILY AS NEEDED
1CQDPM	TAKE 1 CAPSULE BY MOUTH ONCE DAILY IN THE EVENING
1CQDPRN	TAKE 1 CAPSULE BY MOUTH ONCE DAILY AS NEEDED
1CQDQAM	TAKE 1 CAPSULE BY MOUTH ONCE DAILY EVERY MORNING
1CQHS	TAKE 1 CAPSULE BY MOUTH EVERY NIGHT AT BEDTIME
1CQID	TAKE 1 CAPSULE BY MOUTH 4 TIMES A DAY
1CQIDPRN	TAKE 1 CAPSULE BY MOUTH 4 TIMES A DAY AS NEEDED
1CQOD	TAKE 1 CAPSULE BY MOUTH EVERY OTHER DAY
1CQW	TAKE 1 CAPSULE BY MOUTH EVERY WEEK
1CTID	TAKE 1 CAPSULE BY MOUTH 3 TIMES A DAY
1CTIDCF	TAKE 1 CAPSULE BY MOUTH 3 TIMES A DAY WITH FOOD
1CTIDCFPRN	TAKE 1 CAPSULE BY MOUTH 3 TIMES A DAY WITH FOOD AS NEEDED
1CTIDPRN	TAKE 1 CAPSULE BY MOUTH 3 TIMES A DAY AS NEEDED
1SSTHS	TAKE 1 AND 1/2 TABLETS BY MOUTH AT BEDTIME
1TAC	TAKE 1 TABLET BY MOUTH BEFORE MEALS
1TACHS	TAKE 1 TABLET BY MOUTH BEFORE MEALS AND AT BEDTIME
1TACTID	TAKE 1 TABLET BY MOUTH BEFORE MEALS 3 TIMES A DAY
1TACPRN	TAKE 1 TABLET BY MOUTH BEFORE MEALS AS NEEDED
1TAM	TAKE 1 TABLET BY MOUTH IN THE MORNING
1TAM.5TPM	TAKE 1 TABLET BY MOUTH IN THE MORNING AND TAKE 1/2 TABLET BY MOUTH IN THE EVENING

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
1TAM1.5THS	TAKE 1 TABLET BY MOUTH IN THE MORNING AND TAKE 1 AND 1/2 TABLETS BY MOUTH AT BEDTIME
1TAM1TPM	TAKE 1 TABLET BY MOUTH IN THE MORNING AND TAKE 1 TABLET BY MOUTH IN THE EVENING
1TAM2THS	TAKE 1 TABLET BY MOUTH IN THE MORNING AND TAKE 2 TABLETS BY MOUTH AT BEDTIME
1TAM2TPM	TAKE 1 TABLET BY MOUTH IN THE MORNING AND TAKE 2 TABLETS BY MOUTH IN THE EVENING
1TAMHS	TAKE 1 TABLET BY MOUTH IN THE MORNING AND AT BEDTIME
1TAMPM	TAKE 1 TABLET BY MOUTH IN THE MORNING AND IN THE EVENING
1TAMPRN	TAKE 1 TABLET BY MOUTH IN THE MORNING AS NEEDED
1TAMQD	TAKE 1 TABLET BY MOUTH IN THE MORNING ONCE DAILY
1TBID	TAKE 1 TABLET BY MOUTH TWICE DAILY
1TBIDAC	TAKE 1 TABLET BY MOUTH TWICE DAILY BEFORE MEALS
1TBIDCF	TAKE 1 TABLET BY MOUTH TWICE DAILY WITH FOOD
1TBIDPC	TAKE 1 TABLET BY MOUTH TWICE DAILY AFTER MEALS
1TBIDPRN	TAKE 1 TABLET BY MOUTH TWICE DAILY AS NEEDED
1TB-TID	TAKE 1 TABLET BY MOUTH 2 TO 3 TIMES A DAY
1THS	TAKE 1 TABLET BY MOUTH AT BEDTIME
1THSPRN	TAKE 1 TABLET BY MOUTH AT BEDTIME AS NEEDED
1TPM	TAKE 1 TABLET BY MOUTH IN THE EVENING
1TPMCF	TAKE 1 TABLET BY MOUTH IN THE EVENING WITH FOOD
1TPMPRN	TAKE 1 TABLET BY MOUTH IN THE EVENING AS NEEDED
1TPOBID	TAKE 1 TABLET BY MOUTH TWICE DAILY
1TPOBIDPRN	TAKE 1 TABLET BY MOUTH TWICE DAILY AS NEEDED
1TPOHS	TAKE 1 TABLET BY MOUTH AT BEDTIME
1TPOHSPRN	TAKE 1 TABLET BY MOUTH AT BEDTIME AS NEEDED
1TPOQAM	TAKE 1 TABLET BY MOUTH EVERY MORNING
1TPOQD	TAKE 1 TABLET BY MOUTH ONCE DAILY

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
1TPOQDPRN	TAKE 1 TABLET BY MOUTH ONCE DAILY AS NEEDED
1TPOQHS	TAKE 1 TABLET BY MOUTH EVERY DAY AT BEDTIME
1TPOQID	TAKE 1 TABLET BY MOUTH 4 TIMES A DAY
1TPOQIDPRN	TAKE 1 TABLET BY MOUTH 4 TIMES A DAY AS NEEDED
1TPOQM	TAKE 1 TABLET BY MOUTH ONCE MONTHLY
1TPOQOD	TAKE 1 TABLET BY MOUTH EVERY OTHER DAY
1TPOQPM	TAKE 1 TABLET BY MOUTH EVERY EVENING
1TPOQWK	TAKE 1 TABLET BY MOUTH ONCE WEEKLY
1TPOTID	TAKE 1 TABLET BY MOUTH 3 TIMES A DAY
1TPOTIDPRN	TAKE 1 TABLET BY MOUTH 3 TIMES A DAY AS NEEDED
1TQ12H	TAKE 1 TABLET BY MOUTH EVERY 12 HOURS
1TQ2-4	TAKE 1 TABLET BY MOUTH EVERY 2 TO 4 HOURS
1TQ3	TAKE 1 TABLET BY MOUTH EVERY 3 HOURS
1TQ3-4	TAKE 1 TABLET BY MOUTH EVERY 3 TO 4 HOURS
1TQ3-4PRN	TAKE 1 TABLET BY MOUTH EVERY 3 TO 4 HOURS AS NEEDED
1TQ3PRN	TAKE 1 TABLET BY MOUTH EVERY 3 HOURS AS NEEDED
1TQ4-6	TAKE 1 TABLET BY MOUTH EVERY 4 TO 6 HOURS
1TQ4-6CF	TAKE 1 TABLET BY MOUTH EVERY 4 TO 6 HOURS WITH FOOD
1TQ4-6PRN	TAKE 1 TABLET BY MOUTH EVERY 4 TO 6 HOURS AS NEEDED
1TQ4-6PRNCF	TAKE 1 TABLET BY MOUTH EVERY 4 TO 6 HOURS AS NEEDED WITH FOOD
1TQ4H	TAKE 1 TABLET BY MOUTH EVERY 4 HOURS
1TQ4PRN	TAKE 1 TABLET BY MOUTH EVERY 4 HOURS AS NEEDED
1TQ6-8	TAKE 1 TABLET BY MOUTH EVERY 6 TO 8 HOURS
1TQ6-8CF	TAKE 1 TABLET BY MOUTH EVERY 6 TO 8 HOURS WITH FOOD
1TQ6-8PRN	TAKE 1 TABLET BY MOUTH EVERY 6 TO 8 HOURS AS NEEDED
1TQ6-8PRNCF	TAKE 1 TABLET BY MOUTH EVERY 6 TO 8 HOURS AS NEEDED WITH FOOD

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
1TQ6CF	TAKE 1 TABLET BY MOUTH EVERY 6 HOURS WITH FOOD
1TQ6H	TAKE 1 TABLET BY MOUTH EVERY 6 HOURS
1TQ6PRN	TAKE 1 TABLET BY MOUTH EVERY 6 HOURS AS NEEDED
1TQ8H	TAKE 1 TABLET BY MOUTH EVERY 8 HOURS
1TQ8PRN	TAKE 1 TABLET BY MOUTH EVERY 8 HOURS AS NEEDED
1TQAM	TAKE 1 TABLET BY MOUTH EVERY MORNING
1TQAM1THS	TAKE 1 TABLET BY MOUTH EVERY MORNING AND TAKE 1 TABLET BY MOUTH AT BEDTIME
1TQAM2PM	TAKE 1 TABLET BY MOUTH EVERY MORNING AND TAKE 2 TABLETS BY MOUTH IN THE EVENING
1TQAMPRN	TAKE 1 TABLET BY MOUTH EVERY MORNING AS NEEDED
1TQ-BID	TAKE 1 TABLET BY MOUTH ONCE OR TWICE DAILY
1TQD	TAKE 1 TABLET BY MOUTH ONCE DAILY
1TQDAC	TAKE 1 TABLET BY MOUTH ONCE DAILY BEFORE A MEAL
1TQDACAM	TAKE 1 TABLET BY MOUTH ONCE DAILY BEFORE THE MORNING MEAL
1TQDAM	TAKE 1 TABLET BY MOUTH ONCE DAILY IN THE MORNING
1TQDAMPRN	TAKE 1 TABLET BY MOUTH ONCE DAILY IN THE MORNING AS NEEDED
1TQDCB	TAKE 1 TABLET BY MOUTH ONCE DAILY AT BREAKFAST
1TQDCF	TAKE 1 TABLET BY MOUTH ONCE DAILY WITH FOOD
1TQDHS	TAKE 1 TABLET BY MOUTH EVERY DAY AT BEDTIME
1TQDPM	TAKE 1 TABLET BY MOUTH ONCE DAILY IN THE EVENING
1TQDPRN	TAKE 1 TABLET BY MOUTH ONCE DAILY AS NEEDED
1TQDQAM	TAKE 1 TABLET BY MOUTH ONCE DAILY EVERY MORNING
1TQHS	TAKE 1 TABLET BY MOUTH EVERY DAY AT BEDTIME
1TQID	TAKE 1 TABLET BY MOUTH 4 TIMES A DAY
1TQOD	TAKE 1 TABLET BY MOUTH EVERY OTHER DAY
1TQPM	TAKE 1 TABLET BY MOUTH EVERY EVENING
1TQWK	TAKE 1 TABLET BY MOUTH ONCE WEEKLY
1TTID	TAKE 1 TABLET BY MOUTH 3 TIMES A DAY
1-1.5TBID	TAKE 1 TO 1 AND 1/2 TABLETS BY MOUTH TWICE DAILY

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
1.5-2TQD	TAKE 1 AND 1/2 TO 2 TABLETS BY MOUTH ONCE DAILY
1.5T	TAKE 1 AND 1/2 TABLETS BY MOUTH
1.5TAM	TAKE 1 AND 1/2 TABLETS BY MOUTH IN THE MORNING
1.5TAMPRN	TAKE 1 AND 1/2 TABLETS BY MOUTH IN THE MORNING AS NEEDED
1.5TBID	TAKE 1 AND 1/2 TABLETS BY MOUTH TWICE DAILY
1.5TD	TAKE 1 AND 1/2 TABLETS BY MOUTH DAILY
1.5THS	TAKE 1 AND 1/2 TABLETS BY MOUTH AT BEDTIME
1.5THSPRN	TAKE 1 AND 1/2 TABLETS BY MOUTH AT BEDTIME AS NEEDED
1.5TPM	TAKE 1 AND 1/2 TABLETS BY MOUTH IN THE EVENING
1.5TQ6	TAKE 1 AND 1/2 TABLETS BY MOUTH EVERY 6 HOURS
1.5TQAM	TAKE 1 AND 1/2 TABLETS BY MOUTH EVERY MORNING
1.5TQD	TAKE 1 AND 1/2 TABLETS BY MOUTH ONCE DAILY
1.5TQDAM	TAKE 1 AND 1/2 TABLETS BY MOUTH ONCE DAILY IN THE MORNING
1.5TQDPRN	TAKE 1 AND 1/2 TABLETS BY MOUTH ONCE DAILY IN THE EVENING
1.5TQHS	TAKE 1 AND 1/2 TABLETS BY MOUTH EVERY NIGHT AT BEDTIME
1.5TQID	TAKE 1 AND 1/2 TABLETS BY MOUTH 4 TIMES A DAY
1.5TTID	TAKE 1 AND 1/2 TABLETS BY MOUTH 3 TIMES A DAY
1-2CHS	TAKE 1 OR 2 CAPSULES BY MOUTH AT BEDTIME
1-2CQD	TAKE 1 OR 2 CAPSULES BY MOUTH ONCE DAILY
1-2CQDPRN	TAKE 1 OR 2 CAPSULES BY MOUTH ONCE DAILY AS NEEDED
1-2CQHS	TAKE 1 OR 2 CAPSULES BY MOUTH AT BEDTIME
1-2TAM	TAKE 1 OR 2 TABLETS BY MOUTH IN THE MORNING
1-2TBID	TAKE 1 OR 2 TABLETS BY MOUTH TWICE DAILY
1-2TBIDPRN	TAKE 1 OR 2 TABLETS BY MOUTH TWICE DAILY AS NEEDED
1-2TDUD	TAKE 1 OR 2 TABLETS BY MOUTH DAILY AS DIRECTED
1-2THS	TAKE 1 OR 2 TABLETS BY MOUTH AT BEDTIME
1-2THSPRN	TAKE 1 OR 2 TABLETS BY MOUTH AT BEDTIME AS NEEDED
1-2TPM	TAKE 1 OR 2 TABLETS BY MOUTH IN THE EVENING
1-2TPOHS	TAKE 1 OR 2 TABLETS BY MOUTH AT BEDTIME

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
1-2TPRN	TAKE 1 OR 2 TABLETS BY MOUTH AS NEEDED
1-2TQ4	TAKE 1 OR 2 TABLETS BY MOUTH EVERY 4 HOURS
1-2TQ4-6	TAKE 1 OR 2 TABLETS BY MOUTH EVERY 4 TO 6 HOURS
1-2TQ4-6PP	TAKE 1 OR 2 TABLETS BY MOUTH EVERY 4 TO 6 HOURS AS NEEDED FOR PAIN
1-2TQ4-6PRN	TAKE 1 OR 2 TABLETS BY MOUTH EVERY 4 TO 6 HOURS AS NEEDED
1-2TQ4PRN	TAKE 1 OR 2 TABLETS BY MOUTH EVERY 4 HOURS AS NEEDED
1-2TQ6	TAKE 1 OR 2 TABLETS BY MOUTH EVERY 6 HOURS
1-2TQ6-8	TAKE 1 OR 2 TABLETS BY MOUTH EVERY 6 TO 8 HOURS
1-2TQ6-8PRN	TAKE 1 OR 2 TABLETS BY MOUTH EVERY 6 TO 8 HOURS AS NEEDED
1-2TQ6PRN	TAKE 1 OR 2 TABLETS BY MOUTH EVERY 6 HOURS AS NEEDED
1-2TQ8	TAKE 1 OR 2 TABLETS BY MOUTH EVERY 8 HOURS
1-2TQ8PRN	TAKE 1 OR 2 TABLETS BY MOUTH EVERY 8 HOURS AS NEEDED
1-2TQAM	TAKE 1 OR 2 TABLETS BY MOUTH EVERY MORNING
1-2TQD	TAKE 1 OR 2 TABLETS BY MOUTH ONCE DAILY
1-2TQDHS	TAKE 1 OR 2 TABLETS BY MOUTH ONCE DAILY AT BEDTIME
1-2TQDPM	TAKE 1 OR 2 TABLETS BY MOUTH ONCE DAILY IN THE EVENING
1-2TQDPRN	TAKE 1 OR 2 TABLETS BY MOUTH ONCE DAILY AS NEEDED
1-2TQDUD	TAKE 1 OR 2 TABLETS BY MOUTH ONCE DAILY AS DIRECTED
1-2TQHS	TAKE 1 TO 2 TABLETS BY MOUTH EVERY NIGHT AT BEDTIME
1-2TQID	TAKE 1 OR 2 TABLETS BY MOUTH 4 TIMES A DAY
1-2TTID	TAKE 1 OR 2 TABLETS BY MOUTH 3 TIMES A DAY
1-3THS	TAKE 1 TO 3 TABLETS BY MOUTH AT BEDTIME
1-3TQDAM	TAKE 1 TO 3 TABLETS BY MOUTH ONCE DAILY IN THE MORNING
2CAM	TAKE 2 CAPSULES BY MOUTH EVERY MORNING
2CBID	TAKE 2 CAPSULES BY MOUTH TWICE DAILY
2CHS	TAKE 2 CAPSULES BY MOUTH AT BEDTIME
2CPM	TAKE 2 CAPSULES BY MOUTH EVERY EVENING
2CPOBID	TAKE 2 CAPSULES BY MOUTH TWICE DAILY

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
2CPOBIDPRN	TAKE 2 CAPSULES BY MOUTH TWICE DAILY AS NEEDED
2CPOQD	TAKE 2 CAPSULES BY MOUTH ONCE DAILY
2CPOQDPRN	TAKE 2 CAPSULES BY MOUTH ONCE DAILY AS NEEDED
2CPOQID	TAKE 2 CAPSULES BY MOUTH 4 TIMES A DAY
2CPOQIDPRN	TAKE 2 CAPSULES BY MOUTH 4 TIMES A DAY AS NEEDED
2CPOTID	TAKE 2 CAPSULES BY MOUTH 3 TIMES A DAY
2CPOTIDPRN	TAKE 2 CAPSULES BY MOUTH 3 TIMES A DAY AS NEEDED
2CQAM	TAKE 2 CAPSULES BY MOUTH EVERY MORNING
2CQD	TAKE 2 CAPSULES BY MOUTH ONCE DAILY
2CQID	TAKE 2 CAPSULES BY MOUTH 4 TIMES A DAY
2CTID	TAKE 2 CAPSULES BY MOUTH 3 TIMES A DAY
2TAM	TAKE 2 TABLETS BY MOUTH EVERY MORNING
2TAM1TAFT	TAKE 2 TABLETS BY MOUTH IN THE MORNING AND 1 TABLET IN THE AFTERNOON
2TAM1TPM	TAKE 2 TABLETS BY MOUTH IN THE MORNING AND 1 TABLET IN THE EVENING
2TBID	TAKE 2 TABLETS BY MOUTH TWICE DAILY
2THS	TAKE 2 TABLETS BY MOUTH AT BEDTIME
2TPM	TAKE 2 TABLETS BY MOUTH EVERY EVENING
2TPOBID	TAKE 2 TABLETS BY MOUTH TWICE DAILY
2TPOBIDPRN	TAKE 2 TABLETS BY MOUTH TWICE DAILY AS NEEDED
2TPOHS	TAKE 2 TABLETS BY MOUTH AT BEDTIME
2TPOQD	TAKE 2 TABLETS BY MOUTH ONCE DAILY
2TPOQDPRN	TAKE 2 TABLETS BY MOUTH ONCE DAILY AS NEEDED
2TPOQID	TAKE 2 TABLETS BY MOUTH 4 TIMES A DAY
2TPOQIDPRN	TAKE 2 TABLETS BY MOUTH 4 TIMES A DAY AS NEEDED
2TPOTID	TAKE 2 TABLETS BY MOUTH 3 TIMES A DAY
2TPOTIDPRN	TAKE 2 TABLETS BY MOUTH 3 TIMES A DAY AS NEEDED
2TQAM	TAKE 2 TABLETS BY MOUTH EVERY MORNING

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
2TQD	TAKE 2 TABLETS BY MOUTH ONCE DAILY
2TQID	TAKE 2 TABLETS BY MOUTH 4 TIMES A DAY
3CAM	TAKE 3 CAPSULES BY MOUTH EVERY MORNING
2TTID	TAKE 2 TABLETS BY MOUTH 3 TIMES A DAY
3CBID	TAKE 3 CAPSULES BY MOUTH TWICE DAILY
3CPM	TAKE 3 CAPSULES BY MOUTH EVERY EVENING
3CQD	TAKE 3 CAPSULES BY MOUTH DAILY
3CQID	TAKE 3 CAPSULES BY MOUTH FOUR TIMES DAILY
3CTID	TAKE 3 CAPSULES BY MOUTH THREE TIMES DAILY
3TAM	TAKE 3 TABLETS BY MOUTH EVERY MORNING
3TBID	TAKE 3 TABLETS BY MOUTH TWICE DAILY
3THS	TAKE 3 TABLETS BY MOUTH AT BEDTIME
3TPM	TAKE 3 TABLETS BY MOUTH EVERY EVENING
3TQD	TAKE 3 TABLETS BY MOUTH DAILY
3TQID	TAKE 3 TABLETS BY MOUTH FOUR TIMES DAILY
3TTID	TAKE 3 TABLETS BY MOUTH THREE TIMES DAILY
4CBID	TAKE 4 CAPSULES BY MOUTH TWICE DAILY
4CQD	TAKE 4 CAPSULES BY MOUTH DAILY
4CQID	TAKE 4 CAPSULES BY MOUTH FOUR TIMES DAILY
4CTID	TAKE 4 CAPSULES BY MOUTH THREE TIMES DAILY
4TBID	TAKE 4 TABLETS BY MOUTH TWICE DAILY
4TQD	TAKE 4 TABLETS BY MOUTH DAILY
4TQID	TAKE 4 TABLETS BY MOUTH FOUR TIMES DAILY
4TTID	TAKE 4 TABLETS BY MOUTH THREE TIMES DAILY
BISPHO	TAKE 1 TABLET BY MOUTH ONCE A WEEK ON AN EMPTY STOMACH WITH A FULL GLASS OF WATER. REMAIN UPRIGHT AND DO NOT EAT OR DRINK FOR 30 MINUTES.
BISPHO60	TAKE 1 TABLET BY MOUTH ONCE A MONTH ON AN EMPTY STOMACH WITH A FULL GLASS OF WATER. REMAIN UPRIGHT AND DO NOT EAT OR DRINK FOR 60 MINUTES

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
C1GBID	CHEW AND SWALLOW 1 GUMMY BY MOUTH TWICE DAILY
C1GQD	CHEW AND SWALLOW 1 GUMMY BY MOUTH ONCE DAILY
C1TAM	CHEW AND SWALLOW 1 TABLET BY MOUTH IN THE MORNING
C1TBID	CHEW AND SWALLOW 1 TABLET BY MOUTH TWICE DAILY
C1THS	CHEW AND SWALLOW 1 TABLET BY MOUTH AT BEDTIME
C1TPM	CHEW AND SWALLOW 1 TABLET BY MOUTH IN THE EVENING
C1TQD	CHEW AND SWALLOW 1 TABLET BY MOUTH ONCE DAILY
C2GBID	CHEW AND SWALLOW 2 GUMMIES BY MOUTH TWICE DAILY
C2GHS	CHEW AND SWALLOW 2 GUMMIES BY MOUTH AT BEDTIME
C2GQAM	CHEW AND SWALLOW 2 GUMMIES BY MOUTH EVERY MORNING
C2GQD	CHEW AND SWALLOW 2 GUMMIES BY MOUTH ONCE DAILY
C2TAM	CHEW AND SWALLOW 2 TABLETS BY MOUTH IN THE MORNING
C2TBID	CHEW AND SWALLOW 2 TABLETS BY MOUTH TWICE DAILY
C2THS	CHEW AND SWALLOW 2 TABLETS BY MOUTH AT BEDTIME
C2TPM	CHEW AND SWALLOW 2 TABLETS BY MOUTH IN THE EVENING
C2TQD	CHEW AND SWALLOW 2 TABLETS BY MOUTH ONCE DAILY
CPPI	TAKE 1 CAPSULE BY MOUTH DAILY 30 MINUTES BEFORE BREAKFAST
DNC	TAKE 4 CAPSULES BY MOUTH 1 HOUR BEFORE DENTAL APPOINTMENT
DNT	TAKE 4 TABLETS BY MOUTH 1 HOUR BEFORE DENTAL APPOINTMENT
GLUTABSIG	CHEW AND SWALLOW 3 TO 4 TABLETS BY MOUTH AS NEEDED FOR LOW BLOOD SUGAR
LABELSIG	TAKE AS DIRECTED BY MOUTH PER PACKAGE LABELING AS NEEDED
NARATRIP	TAKE 1 TABLET BY MOUTH AT ONSET OF HEADACHE. MAY REPEAT ONE TABLET IN 4 HOURS IF NEEDED FOR A MAX DOSE OF 5MG IN 24 HOURS
NITRO	DISSOLVE 1 TABLET UNDER TONGUE EVERY 5 MINUTES AS NEEDED FOR CHEST PAIN, MAX OF 3 TABLETS IN 15 MINUTES AND CALL 911 IF CHEST PAIN IS UNRELIEVED.

English SIG Code Translations

Combination SIG codes - Oral continued

SIG code	Description
NTGSP	USE 1 TO 2 SPRAYS UNDER THE TONGUE EVERY 5 MINUTES AS NEEDED FOR CHEST PAIN (NO MORE THAN 3 SPRAYS IN 15 MINUTES). CALL 911 IF CHEST PAIN PERSISTS.
PEGLAX	MIX 1 CAPFUL (17 GRAMS) WITH 4 TO 8 OUNCES OF WATER AND DRINK BY MOUTH DAILY
RPO	RINSE MOUTH AFTER EACH USE
RPOW	RINSE MOUTH WITH WATER AFTER EACH USE
SOB	SHORTNESS OF BREATH
TNT	ONE TABLET UNDER THE TONGUE AS NEEDED FOR CHEST PAIN
TPPI	TAKE 1 TABLET BY MOUTH DAILY 30 MINUTES BEFORE BREAKFAST
TRIPTANSIG	TAKE 1 TABLET BY MOUTH AT ONSET OF HEADACHE. MAY REPEAT ONE TABLET IN 2 HOURS IF NEEDED FOR A MAX OF 2 TABLETS IN 24 HOURS

Combination SIG codes - Otic

SIG code	Description
1GAUQD	USE 1 DROP IN BOTH EARS ONCE DAILY
1GADQD	USE 1 DROP IN RIGHT EAR ONE TIME A DAY
1GASQD	USE 1 DROP IN LEFT EAR ONE TIME A DAY

Combination SIG codes - Suppository

SIG code	Description
1SUPPR	UNWRAP AND INSERT 1 SUPPOSITORY INTO THE RECTUM
1SUPPV	UNWRAP AND INSERT 1 SUPPOSITORY INTO THE VAGINA
IVB	INSERT 1 APPLICATORFUL VAGINALLY AT BEDTIME
IVS	INSERT 1 SUPPOSITORY VAGINALLY AT BEDTIME

English SIG Code Translations

Combination SIG codes - Topical

SIG code	Description
AAA	APPLY TO AFFECTED AREA
ALAA	APPLY A THIN LAYER TOPICALLY TO THE AFFECTED AREA
ASAA	APPLY A SMALL AMOUNT TOPICALLY TO THE AFFECTED AREA
KETSHAM	APPLY A SMALL AMOUNT OF SHAMPOO TO THE SCALP ONCE OR TWICE WEEKLY. LATHER SCALP WITH SHAMPOO FOR 5 MINUTES BEFORE RINSING.
PATCHSIG	CHANGE AND APPLY 1 PATCH TO CLEAN, DRY SKIN
LIDOPATCH	CHANGE AND APPLY 1 PATCH TO CLEAN, DRY SKIN FOR 12 HOURS ON AND 12 HOURS OFF